9500041

THE UNITED SHAMES OF AMIERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Michigan Agricultural Experiment Station

There has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED, PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE Plant Variety Protection Office, in the applicant(s) indicated in the said copy, and WHEREAS, upon due examination made, the said applicant(s) is (are) adjudged to be entitled to a certificate of plant variety protection under the LAW.

NOW, therefore, this certificate of plant variety protection is to grant unto the said applicant(s) and the successors, heirs or assigns of the said applicant(s) for the term of twenty years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to exclude others from selling the variety, or offering it for sale, or reproducing it, or importing it, or exporting it, or conditioning it for propagation, or cking it for any of the above purposes, or using it in producing a hybrid or different by therefrom, to the extent provided by the Plant Variety Protection Act. In the lates seed of this variety (1) shall be sold by variety name only as a class of certified hall conform to the number of generations specified by the owner of the rights.

WHEAT, COMMON

'Mendon'

In Vertimonn Marrent, I have hereunto set my hand and caused the seal of the Man's Unriety Hrotection Office to be affixed at the City of Washington, D.C. this thirty-first day of May in the year of our Lord one thousand nine hundred and ninety-six.

Attest;

Marsha A. Star Commissioner

Plant Variety Protection Office Agricultural Wherketing Scruice

Surature of Auriculture

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE DIVISION		determine if a plant variety protection	
(INSTRUCTIONS ON REVERSE)	ECTION CERTIFICATE	2421). Information is held confidential	
·	2. TEMPORARY DESIGNATION :		
	C4227	Mendon	
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)	5. PHONE (include area code)		
109 Agriculture Hall Michigan State University	517/355-0123		
		1 Nov. 15, 1994	
6. GENUS AND SPECIES NAME 7. FAMILY NAME	(Botanical)	N - D AM D PM	
	l l	F Filing and Examination Fee:	
	9. DATE OF DETERMINATION	E め、イスケーバ/	
· · · · · · · · · · · · · · · · · · ·	Dec., 1992	Blat a land	
association, etc.)	-	C C C C C C C C C C C C C C C C C C C	
Agricultural Experiment Station	1	1 \$ 3 00 00	
· · · · · · · · · · · · · · · · · · ·		17-30-46	
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERV	E IN THIS APPLICATION AND RECEIVE ALL PA	PERS	
278 Plant and Soil Sciences Building		•	
Michigan State University			
East Lansing, MI 48824	PHONE (include area ands), 517/2	PER OF/E	
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow INS	STRUCTIONS on reverse)	33-9343	
a. A Exhibit A, Origin and Breeding History of the Variety		•	
c. Exhibit C. Objective Description of Variety		-	
d. King Exhibit D. Additional Description of Variety			
 X Seed Sample (2,500 viable untreated seeds). Date Seed Sample mails 	ed to Plant Variety Protection Office		
g. Ki Filing and Examination Fee (\$2,325) made payable to "Treasurer of the	e United States"	,	
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY Plant Variety Protection Act) YES (If "YES," answer items 16 and 17 be			
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE			
LIMITED AS TO NOMBER OF THE PROPERTY OF THE PR		_	
		D KI CERTIFIED	
		<u></u>).	
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE. OR MARKE	TED IN THE U.S. OR OTHER COLINTRIES?	·	
☑ YES (If "YES," GIVE NAMES OF COUNTRIES AND DATES) U.S □ NO	September 1994		
 The applicant(s) declare(s) that a viable sample of basic seeds of this variety will such regulations as may be applicable. 	be furnished with the application and will be rep	lenished upon request in accordance with	
in section 41, and is entitled to protection under the provisions of section 42 of the	of plant variety, and believe(s) that the variety is the Plant Variety Protection Act.	distinct, uniform, and stable as required	
Applicant(s) is (are) informed that false representation herein can jeopardize prote	ection and result in penalties.		
APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE ASPICIATION FOR PLANT VARIETY PROTECTION CERTIFICATE (INSTRUCTIONS DO REPORT ON PROPERTY OF THE CONTROL OF THE CONTR			
-VI-ASI			
Style Klern	SIGNAL MARKETING SERVICE SCHOOL OF THE VARIETY PROTECTION CERTIFICATE REAL STREET PROTECTION CERTIFICATE RECTIONS ON REVERSE 2. TEMPORARY DESIGNATION CERTIFICATE RECTIONS ON REVERSE 2. TEMPORARY DESIGNATION CERTIFICATE RECTION ON REVERSE 3. VARIETY NAME Mendon TOR OFFICIAL USE ONLY POR CA227 Mendon 7. FAMILY NAME (Botanical) POACEAB 9. DATE OF DETERMINATION DEC., 1992 WE FORM OF ORGANIZATION (Corporation, partnership, OR 12. DATE OF INCORPORATION DOT 13. DATE OF INCORPORATION TIVE(S), IF ANY, TO SERVE IN THIS APPLICATION AND RECEIVE ALL PAPERS 28 1111ding PHONE (Include area code): 517/353-9545 AT SUBMITTED (Follow INSTRUCTIONS on reverse) TIVE(S), IF ANY, TO SERVE IN THIS APPLICATION AND RECEIVE ALL PAPERS 28 1111ding PHONE (Include area code): 517/353-9545 AT SUBMITTED (Follow INSTRUCTIONS on reverse) TO Tale Seed Sample mailed to Plant Variety Protection Office Dayable to Treasure of the United States 17. IF "YES" TO ITEM Is, WHICH CLASSES OF PRODUCTION BEYOND BREEDER SEED? 17. IF "YES" TO ITEM Is, WHICH CLASSES OF PRODUCTION BRYOND BREEDER SEED? 18. THE VARIETY DE SCLD BY VARIETY NAME (N.Y AS A CLASS OF CERTIFIED SEED? (See section 83(a) of the naswer items 16 and 17 below) 19. Date Seed Sample mailed to Plant Variety Protection Office Dayable to "Treasure of the United States" 19. FOUNDATION REGISTERED (SEED) 10. FOUNDATION REGISTERED (SEED) 10. FOUNDATION REGISTERED (SEED) 10. DATE 11. IF "YES" TO ITEM Is, WHICH CLASSES OF PRODUCTION BEYOND BREEDER SEED? 11. If SECURITY IN THE U.S.? 12. POUNDATION REGISTERED (SEED) 13. JOHN RECTION OF THE VARIETY IN THE U.S.? 14. POUNDATION REGISTERED (SEED) 15. POUNDATION REGISTERED (SEED) 16. SEEDER SEED SEED? 17. FOUNDATION REGISTERED (SEED) 18. SECURITY REGISTERED (SEED) 19. DATE 19. DATE 19. DATE 19. DATE 20. DATE 21. SEPTON OF THE VARIETY IN THE U.S.? 22. ORGANICA AND RECEIVED THE U.S.? 23. DATE 24. SEPTON OF THE U.S. THE U.S.? 25. ORGANICA AND RECEIVED THE U.S.? 26. DATE 27. DATE 28. DATE 29. DATE 29.		
GNATURE OF APPLICANT [Owner(s)]	CAPACITY OR TITLE	DATE	
Million of the second of the s			

9500041

Exhibit A. Origin and Breeding History of Mendon Wheat

'Mendon' is a soft red winter wheat (*Triticum aestivum* L.) developed by Michigan State University. Certified seed will be available to farmers in 1994. Mendon was developed primarily for the wheat production regions of southern Michigan.

Mendon is a pure-line selection from the 1978 cross 780756 which has the parentage X0467/B2141//B5250.

The parentage of MSU line X0467 is 'Genesee'/'Winoka'.

The parentage of MSU line B2141 is ('Suwon 92'/'Brevor'//5*'Genesee', A6506)/4/(A4528,Norin 10/Brevor//Yorkwin/3/3*Genesee)

The parentage of MSU line B5250 is (Talbot/CI8487, B0049)/3/(A6200, Genesee*4//Norin 10/Brevor)

 $F_{2:3}$ head rows from the 780756 cross were planted in 1979. $F_{2:4}$ bulk selections were subjected to a further round of head selection resulting in $F_{4:5}$ head rows. Two subsequent generations of bulk selection resulting in an $F_{4:6}$ line designated C4227. That line, which was segregating for grain color, was evaluated in yield trials from 1985 through 1987. Manual separation and increase of white and red seed of C4227 during 1988 generated two sister lines designated C4227 (red seed) and C4827 (white seed). C4227 was evaluated in yield trials in 1989 through 1992. Breeder seed of Mendon was initially constituted in 1990 from yield test derived $F_{4:11}$ seed of C4227. Milling and baking quality evaluations have been conducted on Mendon since 1989.

Mendon has been stable and uniform as described in exhibit "C" in the four years (1990-1993) since creation of the initial lot of breeders seed. Mendon exhibits variants at or below the levels indicated below:

- 1) taller plants, awnless heads, white glume 2.8 percent
- 2) taller plants, awnless head, brown glume 0.9 percent
- 3) taller plants, awned heads, white glume 0.7 percent
- 4) taller plants, awned head, brown glume 0.3 percent
- 5) up to 0.60% white-seeded variant in harvested seed

MICHIGAN STATE UNIVERSITY

WHEAT BREEDING AND GENETICS

Department of Crop and Soil Sciences Room A382 Plant and Soil Sciences Building East Lansing, Michigan 48824-1325 Office (517) 355-2231 / FAX (517) 353-3995 Internet 22857mgr@ibm.cl.msu.edu Richard W. Ward, Associate Professor

2/16/96

Mendon is most similar to Augusta. Both Mendon and Augusta have white chaff at harvest. Mendon flowers earlier than Augusta. In single location yield tests conducted in Ingham county during 1994 and 1995, Mendon reached anthesis an average of 4.3 days earlier than Augusta. That difference was significant at in each of the trials at an alpha level of 0.01 according to LSD values derived from ANOVA analysis. Mendon's kernel color is red while Augusta's is white. On a Munsel color chart, Mendon kernels in samples from 1991 and 1992 yield tests range in color (on the 1975 Munsell Soil Color Chart) from 7.5YR 7/6 to 7.5YR 6/6, while Augusta's were uniformly 10YR 8/4. Mendon's thousand kernel weight (TKW) is larger than Augusta's. In an ANOVA analysis of mean TKW at 19 sites in Michigan (four years), Mendon's TKW was significantly larger than Augusta's (39.5 g Vs. 35.4 g, alpha=0.05, based on the Duncan's multiple range test which was derived from ANOVA analysis of a set of varieties). There is no evidence that error variances were associated with year of trial. Therefore the across environment analysis (which involves averaging data from sites in different years) is most appropriate.



U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK AND SEED DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (TRITICUM SPP.)

	TICUM SPP.)
NAME OF APPLICANT(S)	FOR OFFICIAL USE ONLY
Michigan Agricultural Experiment Station ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	PVPO NUMBER
•	9500041
109 Agriculture Hall	DESIGNATION
Michigan State University	
East Lansing, MI 48824-1039	Mendon
Place the appropriate number that describes the varietal characte Place a zero in first box (e-s. 0 8 9 or 0 9) when number	r of this variety in the boxes below. is either 99 or less or 9 or less.
1. KIND:	
	= POLISH 6 = POULARD 7 = CLUB
2. TYPE,	1 = SOFT 3 = OTHER (Specify)
2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	1 2 = HARD
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	·
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	<u></u>
1 5 9 FIRST FLOWERING	1 6 3 LAST FLOWERING
4. MATURITY (50% Flowering):	
NO. OF DAYS EARLIER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
0 2 NO. OF DAYS LATER THAN	1 4 = LEMHI 5 = NUGAINES 6 = LEEDS
5. PLANT HEIGHT (From soil level to top of head):	
1 1 0 cm. HIGH	
1 5 CM. TALLER THAN	1
CM. SHORTER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS 4 = LEMHI 5 = NUGAINES 6 = LEEDS
6. PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR:
2 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 1 = YELLOW 2 = PURPLE
8. STEM:	
1 Anthocyanin: = ABSENT 2 = PRESENT	1 Waxy bloom: 1 = ABSENT 2 = PRESENT
Hairiness of last internode of rachis: ABSENT 2 = PRESENT	Internodes: 1 = HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground)	2 1 CM. INTERNODE LENGTH BETWEEN FLAG LEAF
AURICLES:	
Anthocyanin: 1 = ABSENT 2 = PRESENT	Hairiness: I = ABSENT 2 = PRESENT
O. LEAF:	
2 Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):	2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED
1 Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT	1 Waxy bloom of flag leaf sheath: I = ABSENT 2 = PRESENT
1 1 MM. LEAF WIDTH (First leaf below flag lead)	2 7 CM. LEAF LENGTH (First leaf below flag leaf):
0.00 1.000 470.0 (0.00) (5	

11. HEAD:			
Density: = LAX	2 = DENSE	1 Shape: 1 = TAPE 4 = OTHE	ERING 2 = STRAP 3 = CLAVATE ER (Specify)
2 Awnedness: 1 = A	WNLESS 2 = APICALLY AWNLETED	3 = AWNLETED 4 = AWN	ED ' P
Color at maturity:	1 = WHITE 2 = YELLOW 3 = PINK 7 5 = BROWN 6 = BLACK 7 = OTH	4 = RED IER (Specify):	. v .
0 8 CM. LENGTH	and the second s	1 0 MM. WIDTH	
12. GLUMES AT MATUR 2 Length: 1 = SHORT 3 = LONG	(CA. 7 mm.) 2 = MEDIUM (CA. 8 mm.) (CA. 9 mm.)	2 Width: 1 = NARRO 3 = WIDE (
Shoulder 1 = WAN shape: 4 = SQUA	TING 2 = OBLIQUE 3 = ROUNDED RE 5 = ELEVATED 6 = APICULATE	1 Beak: 1 = ОВТИS	E 2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLOR	₹:	14. SEEDLING ANTHOO	YANIN.
1 1 = WHITE 2 = F	RED 3 = PURPLE	1 1 = ABSENT	· ·
15. JUVENILE PLANT G	OWTH HABIT:		
3 1 = PROSTRATE	2 = SEMI-ERECT 3 = ERE	ст	
16. SEED:			
3 Shape: 1 = OVATE	2 = OVAL 3 = ELLIPTICAL	Cheek: 1 = ROUNI	DED 2 = ANGULAR
1 Brush. 1 = SHORT	2 = MEDIUM 3 = LONG	1 Brush: 1 = NOT C	COLLARED 2 = COLLARED
Phenol reaction (See instructions):	1 = IVORY 2 = FAWN 3 = LT. BROW 4 = BROWN 5 = BLACK		
		•	•
L Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specify)	
0 8 MM. LENGTH	0 3 мм. wіртн	4 0 GM. PER 1000	SEEDS
17. SEED CREASE:			
1 - 1	LESS OF KERNEL 'WINOKA'	Depth: 1 = 20% O	R LESS OF KERNEL 'SCOUT'
	ESS OF KERNEL 'CHRIS'	_ ,	R LESS OF KERNEL "CHRIS"
	AS WIDE AS KERNEL 'LEMHI' ted, 1 = Susceptible, 2 = Resistant)	3 = 50 % 01	R LESS OF KERNEL 'LEMHI'
0 STEM RUST	LEAF RUST	0 STRIPE RUST	0 LOOSE SMUT
2 POWDERY MILDEW	0 BUNT	(Races)	LOOSE 2M01
19. INSECT: (0 = Not Teste	od, 1 = Susceptible, 2 = Resistant)	OTHER (Specity)	
0 SAWFLY	APHID (Bydv.)	O GREEN BUG	0 CEREAL LEAF BEETLE
OTHER (Specify)	HESSIAN FLY		O JOHNSON DESILE
	RACES:	GP^	B C
		DE	,
CHARACTER CHARACTER	TY MOST CLOSELY RESEMBLES THAT S		
Plant tillering	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Leaf size	Lowell Lowell	Seed size	Lowell
Leaf color	Lowell	Seed shape Coleoptile elongation	Lowell
Leaf carriage	Lowell	Seedling pigmentation	Lowell Lowell

INSTRUCTIONS

NERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form

(a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States. Technical

Exhibit D. Additional Description of Variety (Mendon).

Objective Description

Mendon has large red seeds, is beardless with white chaff at harvest, and is slightly shorter than Augusta. Mendon flowers 3-5 days earlier than Augusta. Mendon's winterhardiness is not known to differ from that of Augusta or Frankenmuth. Its resistance to lodging is poor. Mendon exhibits good field resistance to powdery mildew and wheat spindle streak virus, and possesses an unknown major gene for leaf rust resistance. Races that are virulent against this unknown leaf rust resistance have occurred in Michigan. Mendon will exhibit very susceptible reactions when challenged with virulent leaf rust races.

Mendon is uniform and stable and contains variants at levels at or below the levels indicated below:

- 1) Taller plants, awnless heads, white glume 2.8 percent.
- 2) Taller plants, awnless head, brown glume 0.9 percent.
- 3) Taller plants, awned heads, white glume 0.7 percent.
- 4) Taller plants, awned head, brown glume 0.3 percent.

Performance

Agronomic and quality performance data for Michigan tests conducted between 1986 and 1992 are summarized in the attached table. Data for 1985 through 1987 are for the mixed color progenitor of both C4227 and C4827, now "Mendon" and "Lowell", respectively. In multi-location yield tests conducted in Michigan in 1990 through 1992, Mendon averaged 86.3 bushels per acre across the 19 site-years, compared with 76.7 and 75.6 bushels per acre for Augusta and Frankenmuth, respectively. Mendon's test weight is slightly below average and is usually similar to Augusta's. Yearly evaluations at the USDA Soft Wheat Quality Laboratory in Wooster, Ohio have shown that Mendon exhibits superior milling and baking properties.



9500041 2002

Fichigen State University Wheat Breeding Program: Hulti-Year Parformance Summary (All County sites included) Final Report

· ·		Si.	ngle Year Mul Average Yie (Bushels/ac		3	Across Year Averages (bu./acre)		rages	1992		COLOR :		Anthesis	1992	ōbs.
Variety Name	MSU Stock IO	1989	1990	1991	1992	2 YR 91-92	3 YR 90-92	4 YR 89-92	Test Weight (lbs/bu.)	-	· · · · · · ·	AWNS	Anthesis Date (June 92)	Leaf Rust score	Lodge score
Pioneer 2510	15157	-		_	114.8				59.4	RED	WHITE	OK	6.0	2	ō
Hsu Line	C4826	•	76.4	76.1	108.9	92.5	87.1		55.9	WHITE	8RONZE	NO	5,5	5	3
Kerene	15026		-	66.4	107.3	86.8			57.2	WHITE	WHITE	КО	7.5	2	1
P2737w	15159	-	•		105.7	-		•	55.5	WHITE	WHITE	ЯD	5.5	1	1
Mendon	C4227	-	78.0	76.3	105.0	90.6	86_4		57,9	RED	WHITE	НО	4.5	4	3
Ti486168	14663	+		67.5	104.8	86.1			55.5	WHITE	WHITE	NO	9.0	5	2
Rebecca	15037		•	-	104.8	• •	+	•	57.1	WHITE	WHITE	YES	8.5	7	4
Lowell	C4827		74.9	76.6	103.9	90.2	85.1	•	55.7	WHITE	VHITE	NO	5.5	5	3
Chelsea	C5023	70,5	75,0	66.9	103.9	85.4	81.9	77.0	57.3	WHITE	BRONZE	YES	9.0	3	2
Augusta	M0300	58.6	71.1	61.1	102.8	81.9	78.3	73.4	56.3	WHITE	UHITE	NO	9.0	5	3
Tw86317	14670			64.8	102.2	83,5			55.9	RED	WHITE	KC	8.0	3	2
Harus	M0301	65.9	69.0	71.1	101.7	86.4	80.6	76.9	57.8	WHITE	BRONZE	NO	7.0	2	1
Tu86312	14674	P	-	67.7	100.8	84.2			57.8	RED	WHITE	XO	6.0	3	1
Annette H95-9	15034	•	•	56.9	100.2	78,5	•	•	57.7	WHITE	WHITE	KO	8.5	3	4
Becker	M0297	56.9	61.5	62.1	99.3	80.7	74.3	69.9	57.3	ŘĚD	WHITE	ЖО	4.0	×	1
Cardinal	M0298	62.8	69.5	70.2	98.6	84.4	79.4	75.2	58.6	RED	WHITE	NO	6.5	2	4
Pioneer 2545	15158	•	•		98.2	•			57.7	RED	WHITE	NO	6.0	5	2
Frankermuth	M0290	59.0	66.8	61.9	98.1	60.0	75.6	71.4	58.2	WHITE	BRONZE	NO	9.5	4	4
Hillsdale	M0295	60.2	64.8	60.3	97.2	78.7	74.1	70.6	58.3	RED	BRONZE	NO	8.0	1 .	2
Pioneer 2548	14645	66.7	75.9	75.6	97.2	86.4	82.9	78.8	58_1	RED	WHITE	YES	5.0	2	1
Dynasty	13966	66.9	70.5	62.5	95,7	79.1	76.2	73.9	59.1	RED	UHITE	YES	4.0	4	1
Discovery	15084	•		55.8	95.0	75.4	•		58.8	RED	WHITE	NO	6.D	4	3
Ena	15033	=	•	54.9	94.7	74.8		•	57.4	WHITE	WHITE	YES	8.0	7	3
Ny73116-4u	15072		•	•	94.5	-	-	•	55.8	WHITE	WHITE	МО	9.0	S	Q
Geneva	M0302	63.0	71.7	69.0	94.3	81.6	78.3	74.5	57.8	WHITE	BRONZE	NO	4.5	5	2
Ruby	15156	•	-		92.8	=	-		59.6	RED	WHITE	NO.	5.5	1	· 0
Gr 876	14615	67.7	73.2	71.5	92.6	62.0	79.1	76.2	58.4	RED	WHITE	YES	5.0	1	1
er863	14182	•		-	92.4	•	•	•	57.7	RED	SHITE	YES	2.5	1	х
Pro 100	15160	•	•	•	91.8	-	•		58.4	RED	WHITE	КО	7.5	6	3
Patriot 180	15165	•	-	71.9	91.6	81.7		· ·	611	RED	AHILE	NQ	4.0	4	2
Madison	14631	-	74.4	75.0	91,5	83.2	80.3	-	58.0	RED	WHITE	NO	3.5	2	1
Sauyer	14622	•	•	67.5	91.4	79.4		•	58.0	RED	WHITE	Ю	4.5	1	1
Vakefield	14632	-	79.2	77.0	91.3	84.1	82.5	•	59.6	RED	WHITE	NO	6.5	2	2
Lincoln	14647	61,5	66.0	63.7	90.0	76.8	73.2	70.3	58.6	RED	WHITE	NO	2.5	1	2
Atlantis	15083		-	60.0	89.1	74.5	-	-	60.8	RED	MHITE	NO	4.0	3	3

Intchigan State University Wheat Breeding Program: Multi-Year Performance Summery (All County sites included) Final Report

•	MSU Stock ID	Single Year Hulti-site Average Yields (Bushels/acre)			Acress Year Averages (bu./acre)			19 9 2					1992 Obs.		
Variety None		1989	1990	1991	1992	2 YR 91-92	3 YR 90-92	4 YR 89-92	Test Veight (lbs/bu.)	GRAIN	CHAFF	AVINS	Anthesis Date (June 92)	Leaf Rust score	Lodge score
Teain	14646	68.3	77.3	71.8	88.3	50.0	79-1	76.4	59.8	RED	WHITE	MC	4.5	1	1
Rs 927	15163	•	-	-	88.3	•	-	•	60. 5	RED	WHITE	No.	3.5	1	0
Columbia	15085	•	•	55.3	85.5	70.4			60.1	RED	WHITE	жо	3.5	1	1
\$1203	15071	•		58.1	82.8	70,4	•	•	57.7	RED	WHITE	NO	7.0	6	2
Pro Gold	15162	•		. -	82.8		•		57.6	RED	WHITE	YES	6.0	1	2
Pro 104	15161		•	- ,	81.9	•		•	60.1	RED	WHITE	KQ	3.5	1	2
Patriot 160	15164	•	•	-	81.7		•	•	57.2	RED	WHITE	₩O .	2.5	1	3
Arbor	I5192	•	•		77.1		+		60.0	RED	HITE	110	4.5	1	1
	Hean	63.6	71.9	66.5	95.6	81.6	79.7	74.3	58.0	•					
	Counties	7	7	7 l.e.d.	5 . 8.2 6.9	12	19	26	5 1.2 1.6	÷			:	-	· ·

Notes: 1) Leaf rust and lodging scores based on a 0-9 scale where 0=absent or excellent, and 9= very poor.

2) 1991 and 1992 results primarily from Trial 10 ("State Variety Trial"). For 1990 and 1989, white and red wheats were evaluated in separate Trials which were grown side by side at the various county sites.

3) Data sorted on 1992 everage yield.
For additional information contact R. Ward, 517/355-2231

Exhibit E. Statement of Ownership

The variety for which Plant Variety Protection is hereby sought was developed by Dr. E. Everson and Dr. R. Ward, employees of Michigan State University. All rights to any invention, discovery, or development made by these employees while employed by Michigan State University were assigned to Michigan State University. No rights of any kind pertaining to Mendon are retained by the aforementioned or by any other employees.